

UCSF

UC San Francisco Previously Published Works

Title

Draft Genome Sequence of Mycobacterium arupense Strain GUC1.

Permalink

<https://escholarship.org/uc/item/5rg724nv>

Journal

Genome announcements, 3(3)

ISSN

2169-8287

Authors

Greninger, Alexander L
Cunningham, Gail
Yu, Joanna M
et al.

Publication Date

2015-06-01

DOI

10.1128/genomea.00630-15

Peer reviewed

Draft Genome Sequence of *Mycobacterium arupense* Strain GUC1

Alexander L. Greninger, Gail Cunningham, Joanna M. Yu, Elaine D. Hsu, Charles Y. Chiu, Steve Miller

Department of Laboratory Medicine, UCSF, San Francisco, California, USA

We report the draft genome sequence of *Mycobacterium arupense* strain GUC1 from a sputum sample of a patient with bronchiectasis. This is the first draft genome sequence of *Mycobacterium arupense*, a rapidly growing nonchromogenic mycobacteria.

Received 11 May 2015 Accepted 13 May 2015 Published 11 June 2015

Citation Greninger AL, Cunningham G, Yu JM, Hsu ED, Chiu CY, Miller S. 2015. Draft genome sequence of *Mycobacterium arupense* strain GUC1. *Genome Announc* 3(3):e00630-15. doi:10.1128/genomeA.00630-15.

Copyright © 2015 Greninger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Steve Miller, steve.miller@ucsf.edu.

Mycobacterium arupense is a rapidly growing nonchromogenic mycobacteria that is closely related to the *Mycobacterium terrae* complex and has been isolated from clinical samples, most commonly sputum samples, as well as environmental water sources (1–3). Multiple reports of tenosynovitis and osteoarticular infections with *M. arupense* have also been presented, including infections caused by the type strain AR30097 (4–8). Although the unique identification of *M. arupense* has generally been related to sequence analysis, the phenotypic properties of *M. arupense* that resulted in it being classified as a species include its inability to grow at 42°C, rapid growth at 30°C, variable pyrazinamidase activity, and mycolic acid patterns that distinguish it from *M. terrae* (1).

Rapidly growing mycobacteria constitute a commonly isolated population of acid-fast bacillus in the clinical microbiology lab of varying clinical importance (9, 10). We sequenced the first draft genome of *M. arupense* from a sputum sample of a patient diagnosed with bronchiectasis. The isolate was originally typed as *M. terrae* complex by high-performance liquid chromatography; however, genome sequencing and analysis of the 16S and *rpoB* sequences revealed its identity as *M. arupense*.

DNA from *M. arupense* strain GUC1 was extracted using the Qiagen EZ1 kit, and paired-end libraries were prepared using the Nextera XT DNA library kit followed by sequencing on the Illumina MiSeq. Sequences were adapter and quality (Q20) trimmed using cutadapt, *de novo* assembled using SPAdes v3.5, metagenomically screened for contaminating sequence with SURPI, and annotated via prokka v1.1 (11–14). A total of 6,386,174 paired-end reads of average length 117 nucleotides were recovered after trimming. *De novo* assembly yielded 173 contigs for a total assembly size of 4,441,412 bp with an N_{50} of 56,189 bp, an average coverage of 115×, and a total of 4,182 coding sequences. Contiguity was most likely disrupted by the high G+C content (67%) along with several high-copy-number integrases, transposases, and recombinases that were longer than sequence read length. Other high-copy number contigs included those containing genes to ESX/type VII secretion system, a distantly related 3-methyladenine glycosylase, and a copper-transporting ATPase. The assembly also includes 44 kb across two contigs that aligns with 99 to 100% nucleotide identity to the pMK12478 plasmid from *Mycobacterium kansasii* strain ATCC 12478 (15). Otherwise,

the closest aligning sequenced genomes were *Mycobacterium* sp. JDM601 or *Mycobacterium avium* strains E1/E93 at approximately 80% nucleotide identity. By Comprehensive Antibiotic Resistance Database analysis, the GUC1 strain includes an *ampC* beta-lactamase and two metallo-beta-lactamases which demonstrate 80%, 90%, and 77% amino acid identity to that of *M. avium* strain Env 77, respectively (16, 17).

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. [LASW000000000](https://www.ncbi.nlm.nih.gov/nuclseq/LASW000000000). The assembly described in this paper is the second version, LASW02000000.

ACKNOWLEDGMENT

This work was supported by the University of California San Francisco Department of Laboratory Medicine.

REFERENCES

- Cloud JL, Meyer JJ, Pounder JI, Jost KC, Sweeney A, Carroll KC, Woods GL. 2006. *Mycobacterium arupense* sp. nov., a non-chromogenic bacterium isolated from clinical specimens. *Int J Syst Evol Microbiol* 56:1413–1418. [http://dx.doi.org/10.1099/ijs.0.64194-0](https://doi.org/10.1099/ijs.0.64194-0).
- Liu R, Yu Z, Zhang H, Yang M, Shi B, Liu X. 2012. Diversity of bacteria and mycobacteria in biofilms of two urban drinking water distribution systems. *Can J Microbiol* 58:261–270. [http://dx.doi.org/10.1139/w11-129](https://doi.org/10.1139/w11-129).
- Castillo-Rodal AI, Mazari-Hiriart M, Lloret-Sánchez LT, Sachman-Ruiz B, Vinuesa P, López-Vidal Y. 2012. Potentially pathogenic nontuberculous mycobacteria found in aquatic systems. Analysis from a reclaimed water and water distribution system in Mexico City. *Eur J Clin Microbiol Infect Dis* 31:683–694. [http://dx.doi.org/10.1007/s10096-011-1359-y](https://doi.org/10.1007/s10096-011-1359-y).
- Senda H, Muro H, Terada S. 2011. Flexor tenosynovitis caused by *Mycobacterium arupense*. *J Hand Surg Eur Vol* 36:72–73. [http://dx.doi.org/10.1177/1753193410381825](https://doi.org/10.1177/1753193410381825).
- Seidl A, Lindeque B. 2014. Large joint osteoarticular infection caused by *Mycobacterium arupense*. *Orthopedics* 37:e848–e850. [http://dx.doi.org/10.3928/01477447-20140825-93](https://doi.org/10.3928/01477447-20140825-93).
- Beam E, Vasoo S, Simner PJ, Rizzo M, Mason EL, Walker RC, Deml SM, Brown-Elliott BA, Wallace RJ, Wengenack NL, Sia IG. 2014. *Mycobacterium arupense* flexor tenosynovitis: case report and review of antimicrobial susceptibility profiles for 40 clinical isolates. *J Clin Microbiol* 52:2706–2708. [http://dx.doi.org/10.1128/JCM.00277-14](https://doi.org/10.1128/JCM.00277-14).
- Tsai T, Lai C, Tsai I, Chang C, Hsiao C, Hsueh P. 2008. Tenosynovitis caused by *Mycobacterium arupense* in a patient with diabetes mellitus. *Clin Infect Dis* 47:861–863. [http://dx.doi.org/10.1086/591281](https://doi.org/10.1086/591281).
- Legout L, Ettahar N, Massongo M, Veziris N, Ajana F, Beltrand E, Senneville E. 2012. Osteomyelitis of the wrist caused by *Mycobacterium arupense* in an immunocompetent patient: a unique case. *Int J Infect Dis* 16:e761–e762. [http://dx.doi.org/10.1016/j.ijid.2012.05.007](https://doi.org/10.1016/j.ijid.2012.05.007).

9. Greninger AL, Langelier C, Cunningham G, Keh C, Melgar M, Chiu CY, Miller S. 29 April 2015. Two rapidly growing mycobacterial species isolated from a brain abscess: first whole genome sequence of *Mycobacterium immunogenum* and *Mycobacterium llatzerense*. J Clin Microbiol. <http://dx.doi.org/10.1128/JCM.00402-15>.
10. De Groote MA, Huiitt G. 2006. Infections due to rapidly growing mycobacteria. Clin Infect Dis 42:1756–1763. <http://dx.doi.org/10.1086/504381>.
11. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
12. Naccache SN, Federman S, Veeraraghavan N, Zaharia M, Lee D, Samayoa E, Bouquet J, Greninger AL, Luk K-C, Enge B, Wadford DA, Messenger SL, Genrich GL, Pellegrino K, Grard G, Leroy E, Schneider BS, Fair JN, Martínez MA, Isa P, Crump JA, DeRisi JL, Sittler T, Hackett J, Miller S, Chiu CY. 2014. A cloud-compatible bioinformatics pipeline for ultrarapid pathogen identification from next-generation sequencing of clinical samples. Genome Res 24:1180–1192. <http://dx.doi.org/10.1101/gr.171934.113>.
13. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17:10–12. <http://dx.doi.org/10.14806/ej.17.1.200>.
14. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
15. Wang J, McIntosh F, Radomski N, Dewar K, Simeone R, Enninga J, Brosch R, Rocha EP, Veyrier FJ, Behr MA. 2015. Insights on the emergence of *Mycobacterium tuberculosis* from the analysis of *Mycobacterium kansasii*. Genome Biol Evol 7:856–870. <http://dx.doi.org/10.1093/gbe/evv035>.
16. Hsu C, Wu C, Talaat AM. 2011. Genome-wide sequence variation among *Mycobacterium avium* subspecies *paratuberculosis* isolates: a better understanding of John's disease transmission dynamics. Front Microbiol 2:236. <http://dx.doi.org/10.3389/fmicb.2011.00236>.
17. McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJV, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The comprehensive antibiotic resistance database. Antimicrob Agents Chemother 57:3348–3357. <http://dx.doi.org/10.1128/AAC.00419-13>.